

COMPOUNDS AND METHODS

FIELD OF THE INVENTION

Compounds of this invention are non-peptide, reversible inhibitors of
5 type 2 methionine aminopeptidase, useful in treating conditions mediated by
angiogenesis, such as cancer, haemangioma, proliferative retinopathy,
rheumatoid arthritis, atherosclerotic neovascularization, psoriasis, ocular
neovascularization and obesity.

10 BACKGROUND OF THE INVENTION

In 1974, Folkman proposed that for tumors to grow beyond a critical
size and to spread to form metastases, they must recruit endothelial cells from
the surrounding stroma to form their own endogenous microcirculation in a
process termed angiogenesis (Folkman J. (1974) *Adv Cancer Res.* 19; 331).
15 The new blood vessels induced by tumor cells as their life-line of oxygen and
nutrients also provide exits for cancer cells to spread to other parts of the
body. Inhibition of this process has been shown to effectively stop the
proliferation and metastasis of solid tumors. A drug that specifically inhibits
this process is known as an angiogenesis inhibitor.

20 Having emerged as a promising new strategy for the treatment of
cancer, the anti-angiogenesis therapy ("indirect attack") has several advantages
over the "direct attack" strategies. All the "direct attack" approaches such as
using DNA damaging drugs, antimetabolites, attacking the RAS pathway,
restoring p53, activating death programs, using aggressive T-cells, injecting
25 monoclonal antibodies and inhibiting telomerase, etc., inevitably result in the
selection of resistant tumor cells. Targeting the endothelial compartment of
tumors as in the "indirect attack", however, should avoid the resistance
problem because endothelial cells do not exhibit the same degree of genomic
instability as tumor cells. Moreover, anti-angiogenic therapy generally has
30 low toxicity due to the fact that normal endothelial cells are relatively
quiescent in the body and exhibit an extremely long turnover. Finally since
the "indirect attack" and "direct attack" target different cell types, there is a
great potential for a more effective combination therapy.

More than 300 angiogenesis inhibitors have been discovered, of which
35 about 31 agents are currently being tested in human trials in treatment of
cancers (Thompson, et al., (1999) *J Pathol* 187, 503). TNP-470, a
semisynthetic derivative of fumagillin of *Aspergillus fuigatus*, is among the

most potent inhibitors of angiogenesis. It acts by directly inhibiting endothelial cell growth and migration *in vitro and in vivo* (Ingber et al. (1990) *Nature* 348, 555). Fumagillin and TNP-470, have been shown to inhibit type 2 methionine aminopeptidase (hereinafter MetAP2) by irreversibly modifying its active site. The biochemical activity of fumagillin analogs has been shown to correlate to their inhibitory effect on the proliferation of human umbilical vein endothelial cells (HUVEC). Although the mechanism of the selective action of fumagillin and related compounds on MetAP2-mediated endothelial cell cytostatic effect has not yet been established, possible roles of MetAP2 in cell proliferation have been suggested.

First, hMetAP-2-catalyzed cleavage of the initiator methionine of proteins could be essential for releasing many proteins that, after myristoylation, function as important signaling cellular factors involved in cell proliferation. Proteins known to be myristoylated include the src family tyrosine kinases, the small GTPase ARF, the HIV protein nef and the α subunit of heterotrimeric G proteins. A recently published study has shown that the myristoylation of nitric oxide synthase, a membrane protein involved in cell apoptosis, was blocked by fumagillin (Yoshida, et al. (1998) *Cancer Res.* 58(16), 3751). This is proposed to be an indirect outcome of inhibition of MetAP2-catalyzed release of the glycine-terminal myristoylation substrate. Alternatively, MetAP enzymes are known to be important to the stability of proteins *in vivo* according to the "N-end rule" which suggests increased stability of methionine-cleaved proteins relative to their N-terminal methionine precursors (Varshavsky, A (1996) *Proc. Natl. Acad. Sci. U.S.A.* 93, 12142). Inhibition of hMetAP2 could result in abnormal presence or absence of some cellular proteins critical to the cell cycle.

Methionine aminopeptidases (MetAP) are ubiquitously distributed in all living organisms. They catalyze the removal of the initiator methionine from newly translated polypeptides using divalent metal ions as cofactors. Two distantly related MetAP enzymes, type 1 and type 2, are found in eukaryotes, which at least in yeast, are both required for normal growth; whereas only one single MetAP is found in eubacteria (type 1) and archaeobacteria (type 2). The N-terminal extension region distinguishes the methionine aminopeptidases in eukaryotes from those in procaryotes. A 64-amino acid sequence insertion (from residues 381 to 444 in hMetAP2) in the catalytic C-terminal domain distinguishes the MetAP-2 family from the MetAP-1 family. Despite the difference in the gene structure, all MetAP

enzymes appear to share a highly conserved catalytic scaffold termed "pita-bread" fold (Bazan, et al. (1994) *Proc. Natl. Acad. Sci. U.S.A.* 91, 2473), which contains six strictly conserved residues implicated in the coordination of the metal cofactors.


5 Mammalian type 2 methionine aminopeptidase has been identified as a bifunctional protein implicated by its ability to catalyze the cleavage of N-terminal methionine from nascent polypeptides (Bradshaw, et al (1998) *Trends Biochem. Sci.* 23, 263) and to associate with eukaryotic initiation factor 2 α (eIF-2 α) to prevent its phosphorylation (Ray, et al. (1992) *Proc. Natl. Acad. Sci. U.S.A.* 89, 539). Both the genes of human and rat MetAP2 were cloned and have shown 92% sequence identity (Wu, et al. (1993) *J Biol. Chem.* 268, 10796; Li, X. & Chang, Y.-H. (1996) *Biochem. & Biophys. Res. Comm.* 227, 152). The N-terminal extension in these enzymes is highly charged and consists of two basic polylysine blocks and one aspartic acid block, which has
10 been speculated to be involved in the binding of eIF-2 α (Gupta, et al. (1993) in *Translational Regulation of Gene Expression 2* (Ilan, J., Ed.), pp. 405-431, Plenum Press, New York).

The anti-angiogenic compounds, fumagillin and its analogs, have been shown to specifically block the exo-aminopeptidase activity of hMetAP2
20 without interfering with the formation of the hMetAP2 : eIF2 α complex (Griffith, et al., (1997) *Chem. Biol.* 4, 461; Sin, et al. (1997) *Proc. Natl. Acad. Sci. U.S.A.* 94, 6099). Fumagillin and its analogs inactivate the enzymatic activity of hMetAP2 with a high specificity, which is underscored by the lack of effect of these compounds on the closely related type 1 methionine
25 aminopeptidase (MetAP1) both *in vitro* and *in vivo* in yeast (Griffith, et al., (1997) *Chem. Biol.* 4, 461; Sin, et al. (1997) *Proc. Natl. Acad. Sci. U.S.A.* 94, 6099). The extremely high potency (IC₅₀ < 1 nM) of these inhibitors appears to be due to the irreversible modification of the active site residue, His231, of hMetAP2 (Liu, et al. (1998) *Science* 282, 1324). Disturbance of MetAP2
30 activity *in vivo* impairs the normal growth of yeast (Griffith, et al., (1997) *Chem. Biol.* 4, 461; Sin, et al. (1997) *Proc. Natl. Acad. Sci. U.S.A.* 94, 6099; In-house data) as well as *Drosophila* (Cutforth & Gaul (1999) *Mech. Dev.* 82, 23). Most significantly, there appears to be a clear correlation between the inhibition effect of fumagillin related compounds against the enzymatic
35 activity of hMetAP2 *in vitro* and the suppression effect of these compounds against tumor-induced angiogenesis *in vivo* (Griffith, et al., (1997) *Chem. Biol.* 4, 461).

Cancer is the second leading cause of death in the U.S., exceeded only by heart disease. Despite recent successes in therapy against some forms of neoplastic disease, other forms continue to be refractory to treatment. Thus, cancer remains a leading cause of death and morbidity in the United States and elsewhere (Bailar and Gornik (1997) *N Engl J Med* 336, 1569). Inhibition of hMetAP2 provides a promising mechanism for the development of novel anti-angiogenic agents in the treatment of cancers. It has now been discovered that compounds of formulae (I) and (IA) are effective inhibitors of hMetAP2, and thus would be useful in treating conditions mediated by hMetAP2.

SUMMARY OF THE INVENTION

In one aspect, the present invention is to a compound of formula (I), or a pharmaceutically active salt or solvate thereof, and its use in treating conditions mediated by angiogenesis, such as cancer, haemangioma, proliferative retinopathy, rheumatoid arthritis, atherosclerotic neovascularization, psoriasis, ocular neovascularization and obesity:



Formula (I)

wherein:

X is S or O;


R¹ is optionally substituted C₂₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl;

R² is optionally substituted C₂₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl, provided that when R² is optionally substituted Het-C₀alkyl, and Het is indole, benzofuran, benzothiophene, benzisoxazole, benzothiazole or benzopyrazole, then the optional substituent is not -(CH₂)₂NR⁴R⁵; and

R³ is H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl, C₀₋₆alkyl-C(O)X'AB, C₀₋₆alkyl-

S(O)₂X'AB, C₀₋₆alkyl-X'AB, wherein X' is O, S, C or N; A and B are independently H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl, or A or B are independently
 5 absent, provided that the compound is not 5-anilino-3-benzylthio-1,2,4-triazole, 3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole, 3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole, 3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole, 3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole, or 3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole.

10 In a second aspect, the present invention is to a method of treating conditions mediated by angiogenesis, such as cancer, haemangioma, proliferative retinopathy, rheumatoid arthritis, atherosclerotic neovascularization, psoriasis, ocular neovascularization and obesity by administering a compound of formula (IA), or a pharmaceutically acceptable
 15 salt or solvate thereof



Formula (IA)

wherein,

X is S or O;

20 R¹ is optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl;

R² is optionally substituted C₂₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl;
 25

R³ is H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl, C₀₋₆alkyl-C(O)X'AB, C₀₋₆alkyl-S(O)₂X'AB, C₀₋₆alkyl-X'AB, wherein X' is O, S, C or N; A and B are independently H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted
 30 Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl, or A or B are independently absent.

In another aspect, the present invention is to a method of inhibiting
 35 MetAP2 in the treatment of angiogenesis-mediated diseases, all in mammals,

preferably humans, comprising administering to such mammal in need thereof, a compound of formula (IA), or a pharmaceutically active salt or solvate thereof.

In yet another aspect, the present invention is to pharmaceutical compositions comprising a compound of formula (I) and a pharmaceutically acceptable carrier therefor. In particular, the pharmaceutical compositions of the present invention are used for treating MetAP2-mediated diseases.

DETAILED DESCRIPTION OF THE INVENTION

It has now been discovered that substituted 1,2,4-triazoles of formulae (I) and (IA) are inhibitors of MetAP2. It has also now been discovered that selective inhibition of MetAP2 enzyme mechanisms by treatment with the inhibitors of formula (IA), or a pharmaceutically acceptable salt or solvate thereof, represents a novel therapeutic and preventative approach to the treatment of a variety of disease states; including, but not limited to, cancer, haemangioma, proliferative retinopathy, rheumatoid arthritis, atherosclerotic neovascularization, psoriasis, ocular neovascularization and obesity.

The term "C₁₋₆alkyl" as used herein at all occurrences means a substituted and unsubstituted, straight or branched chain radical of 1 to 6 carbon atoms, unless the chain length is limited thereto, including, but not limited to methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and t-butyl, pentyl, n-pentyl, isopentyl, neopentyl and hexyl and the simple aliphatic isomers thereof. Any C₁₋₆alkyl group may be optionally substituted independently by one or more of OR⁴, R⁴, NR⁴R⁵. C₀alkyl means that no alkyl group is present in the moiety. Thus, Ar-C₀alkyl is equivalent to Ar.

As used herein at all occurrences, substituents R⁴, R⁵, and R⁶ are independently defined as C₂₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, Ar-C₀₋₆alkyl, Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl.

The term "C₃₋₇cycloalkyl" as used herein at all occurrences means substituted or unsubstituted cyclic radicals having 3 to 7 carbons, including but not limited to cyclopropyl, cyclopentyl, cyclohexyl and cycloheptyl radicals.

The term "C₃₋₆alkenyl" as used herein at all occurrences means an alkyl group of 3 to 6 carbons wherein a carbon-carbon single bond is replaced by a carbon-carbon double bond. C₃₋₆alkenyl includes 1-propene, 2-propene, 1-butene, 2-butene, isobutene and the several isomeric pentenes and hexenes. Both cis and trans isomers are included within the scope of this invention.

Any C₃₋₆alkenyl group may be optionally substituted independently by one or more of Ph-C₀₋₆alkyl, Het'-C₀₋₆ alkyl, C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆mercaptyl, Ph-C₀₋₆alkoxy, Het'-C₀₋₆alkoxy, OH, NR⁴R⁵, Het'-S-C₀₋₆alkyl, (CH₂)₁₋₆OH, (CH₂)₁₋₆NR⁴R⁵, O(CH₂)₁₋₆NR⁴R⁵, (CH₂)₀₋₆CO₂R⁶,
 5 O(CH₂)₁₋₆CO₂ R⁶, (CH₂)₁₋₆SO₂, CF₃, OCF₃ or halogen.

The term "C₃₋₆alkynyl" as used herein at all occurrences means an alkyl group of 3 to 6 carbons wherein one carbon-carbon single bond is replaced by a carbon-carbon triple bond. C₃₋₆ alkynyl includes 1-propyne, 2-propyne, 1-butyne, 2-butyne, 3-butyne and the simple isomers of pentyne and
 10 hexyne.

The terms "Ar" or "aryl" as used herein interchangeably at all occurrences mean phenyl and naphthyl, optionally substituted by one or more of Ph-C₀₋₆alkyl, Het'-C₀₋₆ alkyl, C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆mercaptyl, Ph-C₀₋₆alkoxy, Het'-C₀₋₆alkoxy, OH, NR⁴R⁵, Het'-S-C₀₋₆alkyl, (CH₂)₁₋₆OH,
 15 (CH₂)₁₋₆NR⁴R⁵, O(CH₂)₁₋₆NR⁴R⁵, (CH₂)₀₋₆CO₂R⁶, O(CH₂)₁₋₆CO₂ R⁶, (CH₂)₁₋₆SO₂, CF₃, OCF₃ or halogen; in addition, Ph may be optionally substituted with one or more of C₁₋₆alkyl, C₁₋₆alkoxy, OH, (CH₂)₁₋₆NR⁴R⁵, O(CH₂)₁₋₆NR⁴R⁵, CO₂R⁶, CF₃, or halogen; Het' is defined as for Het, and may be optionally substituted by one or more of C₁₋₆alkyl, C₁₋₆alkoxy, OH, (CH₂)₁₋₆NR⁴R⁵, O(CH₂)₁₋₆NR⁴R⁵, CO₂R⁶, CF₃, or halogen;
 20 or two C₁₋₆alkyl or C₁₋₆alkoxy groups may be combined to form a 5-7 membered, saturated or unsaturated ring, fused onto the Ar ring.

Suitably, for compounds of formula (I), when Ar is substituted by Ph or Het', then Ph or Het' are substituted with one or more of C₂₋₆alkyl, C₁₋₆alkoxy, (CH₂)₁₋₆NR⁴R⁵, O(CH₂)₁₋₆NR⁴R⁵, CO₂R⁶, CF₃ or halogen.
 25

The terms "Het" or "heterocyclic" as used herein interchangeably at all occurrences, mean a stable 5- to 7-membered monocyclic, a stable 7- to 10-membered bicyclic, or a stable 11- to 18-membered tricyclic heterocyclic ring, all of which are either saturated or unsaturated, and consist of carbon atoms
 30 and from one to three heteroatoms selected from the group consisting of N, O and S, and wherein the nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The heterocyclic ring may be attached at any
 35 heteroatom or carbon atom which results in the creation of a stable structure.

It will be understood that Het may be optionally substituted with one or more of Ph-C₀₋₆alkyl, Het'-C₀₋₆ alkyl, C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆mercaptyl, Ph-C₀₋₆alkoxy, Het'-C₀₋₆alkoxy, OH, NR⁴R⁵, Het'-S-C₀₋₆alkyl, (CH₂)₁₋₆OH, (CH₂)₁₋₆NR⁴R⁵, O(CH₂)₁₋₆NR⁴R⁵, (CH₂)₀₋₆CO₂R⁶,
 5 O(CH₂)₁₋₆CO₂ R⁶, (CH₂)₁₋₆SO₂, CF₃, OCF₃, CN, or halogen; Ph may be optionally substituted with one or more of C₁₋₆alkyl, C₁₋₆alkoxy, OH, (CH₂)₁₋₆NR⁴R⁵, O(CH₂)₁₋₆NR⁴R⁵, CO₂R⁶, CF₃, or halogen; and two C₁₋₆alkyl or C₁₋₆alkoxy groups may be combined to form a 5-7 membered ring, saturated or unsaturated, fused onto the Het ring. Preferred optional
 10 substituents on Het are C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆mercaptyl, halogen, CF₃, OCF₃, CN, or NR⁴R⁵.

Het' is defined as for Het and may be optionally substituted by one or more of C₁₋₆alkyl, C₁₋₆alkoxy, OH, (CH₂)₁₋₆NR⁴R⁵, O(CH₂)₁₋₆NR⁴R⁵, CO₂R⁶, CF₃, or halogen.

15 Examples of such heterocycles include, but are not limited to piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolodinyl, 2-oxoazepinyl, azepinyl, pyrrolyl, 4-piperidinyl, pyrrolidinyl, pyrazolyl, pyrazolidinyl, imidazolyl, pyridinyl, pyrazinyl, oxazolidinyl, oxazolinyl, oxazolyl, isoxazolyl, morpholinyl, thiazolidinyl,
 20 thiazolinyl, thiazolyl, quinuclidinyl, indolyl, quinolinyl, isoquinolinyl, benzimidazolyl, benzopyranyl, benzoxazolyl, furyl, pyranyl, tetrahydrofuryl, tetrahydropyranyl, thienyl, benzoxazolyl, benzofuranyl, benzothiophenyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, and oxadiazolyl, as well as triazolyl, thiadiazolyl, oxadiazolyl, isoxazolyl, isothiazolyl, imidazolyl,
 25 pyridazinyl, pyrimidinyl and triazinyl which are available by routine chemical synthesis and are stable.

Compounds of this invention of formula (I), do not include compounds wherein R² is optionally substituted Het-C₀alkyl, and Het is indole, benzofuran, benzothiophene, benzisoxazole, benzothiozole or benzopyrazole,
 30 and the optional substituent is -(CH₂)₂NR⁴R⁵. The following compounds of this invention are known: 3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole, 3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole, 3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole, 3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole, or 3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole. Fromm et al., *Justus Liebigs Ann. Chem.*, 437 1924, 113. A compound of formula (I) wherein R¹ is benzyl,
 35 R² is phenyl and R³ is hydrogen is known.

Suitably, when moieties R¹, R², or R³ are either optionally substituted Ar-C₀₋₆alkyl or optionally substituted Het-C₀₋₆alkyl, the moiety may be attached to the triazole substituent through the aromatic ring or through the alkyl chain.

5 Further, it will be understood that when a moiety is "optionally substituted" the moiety may have one or more optional substituents, each optional substituent being independently selected.

The terms "hetero" or "heteroatom" as used herein interchangeably at all occurrences mean oxygen, nitrogen and sulfur.

10 The terms "halo" or "halogen" as used herein interchangeably at all occurrences mean F, Cl, Br, and I.

Here and throughout this application the term C₀ denotes the absence of the substituent group immediately following; for instance, in the moiety ArC₀₋₆alkyl, when C is 0, the substituent is Ar, e.g., phenyl. Conversely,
15 when the moiety ArC₀₋₆alkyl is identified as a specific aromatic group, e.g., phenyl, it is understood that C is 0.

Suitably X is sulfur or oxygen. Preferably X is sulfur.


Suitably, R¹ is optionally substituted C₂₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl. Preferably R¹ is optionally substituted
20 Ar-C₀₋₆alkyl or optionally substituted Het-C₀₋₆alkyl. More preferably R¹ is optionally substituted Ar-C₁alkyl or optionally substituted Het-C₁alkyl. Most preferably R¹ is optionally substituted benzyl, optionally substituted methylfuran or optionally substituted methylthiophene. Preferably, when R¹
25 is Het-C₁alkyl, the alkyl chain is directly attached to moiety X.

Suitably, R² is optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl. Preferably, R² is optionally substituted Ar-C₀₋₆alkyl. More preferably R² is optionally substituted Ar-C₀alkyl. Most
30 preferably R² is optionally substituted Ar-C₀alkyl, wherein the optional substituent is ortho C₁₋₆alkyl, preferably branched C₁₋₆alkyl, most preferably isopropyl.

Suitably, R³ is H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl, C₀₋₆alkyl-C(O)X'AB, C₀₋₆alkyl-S(O)₂X'AB, C₀₋₆alkyl-X'AB, wherein X' is O, S, C or N; A and B are
35 independently H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl,

optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl, or A or B are independently absent. Preferably R³ is hydrogen or C₀₋₆alkyl-C(O)X'AB. More preferably R³ is hydrogen or C₀₋₆alkyl-C(O)X'AB, wherein X' is oxygen and A is methyl or hydrogen and B is absent.

A preferred compound of this invention is a compound of formula (IB):



Formula (IB).

Suitably, pharmaceutically acceptable salts of formula (I) include, but are not limited to, salts with inorganic acids such as hydrochloride, sulfate, phosphate, diphosphate, hydrobromide, and nitrate, or salts with an organic acid such as malate, maleate, fumarate, tartrate, succinate, citrate, acetate, lactate, methanesulfonate, p-toluenesulfonate, palmitate, salicylate, and stearate.

The compounds of the present invention may contain one or more asymmetric carbon atoms and may exist in racemic and optically active forms. The stereocenters may be (R), (S) or any combination of R and S configuration, for example, (R,R), (R,S), (S,S) or (S,R). All of these compounds are within the scope of the present invention.

All compounds of formula (IA) specifically named herein are considered to be part of the invention disclosed herein. Among the compounds of the invention of formula (IA) are the following compounds:

- 3-anilino-5-benzylthio-1,2,4-triazole;
- 3-anilino-5-methylthio-1,2,4-triazole;
- 3-anilino-5-(4-chloro-benzylthio)-1,2,4-triazole;
- 3-anilino-5-allylthio-1,2,4-triazole;
- 3-anilino-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
- 3-anilino-5-(2-methyl-butylthio)-1,2,4-triazole;
- 3-anilino-5-(2-methyl-2-pentenylthio)-1,2,4-triazole;
- 3-anilino-5-(α -methylbenzylthio)-1,2,4-triazole;
- 3-anilino-5-(cyclohexylmethylthio)-1,2,4-triazole;
- 3-anilino-5-(propyl acetylthio)-1,2,4-triazole;

- 3-anilino-5-(3,3-dimethoxy-propylthio)-1,2,4-triazole;
3-anilino-5-(2-phenethylthio)-1,2,4-triazole;
3-anilino-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(3-phenyl-[1,2,4]oxadiazol-5-ylmethylthio)-1,2,4-triazole;
5 3-anilino-5-(1*H*-benzoimidazol-2-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(2-(4-chlorophenyl)-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(4-*i*-propyl-benzylthio)-1,2,4-triazole;
10 3-anilino-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(quinolin-8-ylthio)-1,2,4-triazole;
3-anilino-5-(4-acetamido-benzylthio)-1,2,4-triazole;
4-(5-anilino-2 *H*-[1,2,4]triazol-3-yl thio)-benzoic acid;
3-anilino-5-(2-methyl-benzylthio)-1,2,4-triazole;
15 3-anilino-5-(4-trifluoromethyl-benzylthio)-1,2,4-triazole;
3-anilino-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-anilino-5-(3,5-dimethyl-benzylthio)-1,2,4-triazole;
3-anilino-5-(4-cyano-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
20 3-(4-methyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
25 3-(4-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
30 3-(4-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
35 3-(2-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;

- 3-(2-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
 3-(2-methyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
 3-(2-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
 5 3-(4-chloro-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
 10 3-(4-chloro-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
 15 3-(4-methoxy-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
 20 3-(4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
 4-(5-(cyclohexylmethylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid
 25 methyl ester;
 4-(5-(pyridin-4-ylmethylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid
 methyl ester;
 4-(5-(2-methyl-2-butenylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid
 methyl ester;
 30 4-(5-(2-fluoro-benzylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl
 ester;
 4-(5-(5-methyl-isoxazol-3-ylmethylthio)-1H-[1,2,4]triazol-3-ylamino)-
 benzoic acid methyl ester;
 4-(5-(3-methoxy-benzylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid
 35 methyl ester;
 4-(5-(2-methyl-benzylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl
 ester;

- 4-(5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
- 4-(5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
- 5 4-(5-(2-methyl-thiazol-4-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
- 4-(5-(pyridin-2-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
- 3-(3,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
- 10 3-(3,4-dimethoxy-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole;
- 3-(3,4-dimethoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
- 3-(3,4-dimethoxy-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
- 3-(3,4-dimethoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
- 3-(3,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
- 15 3-(3,4-dimethoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
- 3-(3,4-dimethoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
- 3-(3,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
- 3-(3,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
- 20 3-(3,4-dimethoxy-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
- 3-(3,4-dimethoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-benzylthio-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole;
- 25 3-(2-phenyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
- 30 3-(2-phenyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
- [5-(benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
- 35 [5-(3-methoxybenzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
- [5-(cyclohexylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
- [5-(pyridin-4-ylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;

[5-(2-methyl-2-butenylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-fluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(5-methyl-isoxazol-3-ylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;

- 5 [5-(2-methyl-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(pyridin-2-ylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methyl-thiazol-4-ylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
- 10 [5-(thiophen-2-ylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
3-(2-ethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
- 15 3-(2-ethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
- 20 3-(2-ethyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
- 25 3-(2-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
- 30 3-(2-methoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
- 35 3-(2-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;

- 3-(2-isopropyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
5 3-(2-isopropyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
10 3-(2-isopropyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
15 3-(3-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
20 3-(3-methyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
25 3-(4-*n*-butyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
30 3-(4-*n*-butyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
35 3-(2,4-dimethoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;

- 3-(2,4-dimethoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
5 3-(2,4-dimethoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
10 3-(2,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
15 3-(2-methyl-4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
20 3-(2-methyl-4-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
25 3-(2,6-dimethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
30 3-(2,6-dimethyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-methyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-*n*-propyl-3-anilino-5-benzylthio-1,2,4-triazole;
35 3-*n*-butyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-*i*-propyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-allyl-3-anilino-5-benzylthio-1,2,4-triazole; and

3-benzyl-3-anilino-5-benzylthio-1,2,4-triazole.

Among the preferred compounds of formula (IA) of this invention are the following compounds:

- 5 3-anilino-5-benzylthio-1,2,4-triazole;
3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
- 10 3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-(4-chloro-anilino)-5-benzylthio-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-ethyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-benzylthio-1,2,4-triazole;
- 15 3-(3-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-benzylthio-1,2,4-triazole;
- 20 3-methylacetate-3-(*p*-methyl)-anilino-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-(*p*-methoxy)-anilino-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-(2,6-dimethyl)-anilino-5-benzylthio-1,2,4-triazole;
3-anilino-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
- 25 5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(4-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
- 30 3-(4-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
- 35 3-(4-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;

- 3-(2-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
5 3-(2-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic
acid ethyl ester;
3-(2-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-
10 carbaldehyde;
3-(2-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
15 [5-(thiophen-2-ylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
3-(2-ethyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
5-(5-(2-methoxyphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-
20 carboxylic acid ethyl ester
3-(2-methoxy-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
25 3-(3-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-
30 carboxylic acid ethyl ester;
3-(3-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-
carbaldehyde;
3-(4-*n*-butyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
35 3-(2,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;

- 3-(4-fluoro-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5 5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(4-fluoro-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
10 3-anilino-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
15 3-(2-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
20 4-(5-(3-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
4-(5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
4-(5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid
25 methyl ester;
3-(3,4-dimethoxy-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
30 [5-(2-fluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(thiophen-2-ylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
3-(2-ethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
35 3-(2-ethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;

- 3-(2-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
5 3-(3-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
10 3-(2,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
15 3-(2,6-dimethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole; and
3-(4-fluoro-anilino)-5-(furan-3-ylthio)-1,2,4-triazole.

20

Among the more preferred compounds of formula (IA) are the following compounds:

- 3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole;
25 3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-anilino-5-benzylthio-1,2,4-triazole;
4-(5-benzylthio-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
3-anilino-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(furan-3-ylthio)-1,2,4-triazole;
30 3-anilino-5-(furan-2-ylthio)-1,2,4-triazole;
3-anilino-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-anilino-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
35 3-(4-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;


- 3-(2-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
5 3-(2-methoxy-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole; and
3-(3-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole.

- 10 Among the most preferred compounds of formula (IA) are the following compounds:
3-(2-isopropyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
15 3-(2-isopropyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(2-isopropylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
20 5-(5-(2-isopropyl amino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(2-isopropyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
25 3-(4-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole; and
3-(2-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole.

Methods of Preparation

- 30 Compounds of the formulae (I) and (IA) wherein X is S and R³ is H, were prepared by methods analogous to those described in Scheme 1.

Scheme 1




- 5 a) Thiourea, NaOH, H₂O/CH₃CN; b) EtI, Et₃N, DMF; c) H₂NNH₂, EtOH; d)
 R¹X (X = halogen), K₂CO₃, DMF; e) ClCH₂OCH₂CH₃, NaH, THF; f)
 R³CH₂Br, NaH, DMF; g) TFA.

An isothiocyanate (such as phenyl isothiocyanate) (1-Scheme 1) was treated with thiourea and sodium hydroxide in acetonitrile/water to provide 2-Scheme 1, which was treated with iodoethane and triethylamine in DMF to afford 3-Scheme 1. Treatment of 3-Scheme 1 with hydrazine in ethanol provided 4-Scheme 1, which was treated with an alkyl halide (such as benzyl bromide or 4-chlorobenzyl chloride) and potassium carbonate in DMF to give 5-Scheme 1. Triazole 5-Scheme 1 is protected as the methoxy methylethyl ether to afford 6-Scheme 1. Alkylation of 6-Scheme 1 with an alkyl halide (such as methyl iodide, ethyl iodide, *i*-isobutyl iodide, *n*-propyl iodide, butyl iodide, allyl bromide, benzyl bromide, and methyl bromoacetate) afforded the desired tertiary amine 7-Scheme 1. Deprotection of the MOM-ether 7-Scheme 1 with trifluoroacetic acid (TFA) provided the desired product 8-Scheme 1.

Compounds of the formulae (I) and (IA) wherein X is O may be prepared by methods analogous to those described in Scheme 2.

Scheme 2



- a) Thiourea, EtI, EtOH; b) NH_2NH_2 , EtOH c) 1,1'-Carbonyldiimidazole, EtOH; d) R^1X (X = halogen), K_2CO_3 , DMF; e) $\text{ClCH}_2\text{OCH}_2\text{CH}_3$, NaH, THF; f) $\text{R}^3\text{CH}_2\text{Br}$, NaH, DMF; g) TFA.

A thiourea (such as phenylthiourea) (8-Scheme 2) may be treated with ethyl iodide and refluxed in EtOH, and the resulting S-ethyl thiourea is then heated in the presence of hydrazine to provide 9-Scheme 2. The hydrazine 9-Scheme 2 is treated with carbonyldiimidazole and heated to afford 10-Scheme 2. Treatment of 10-Scheme 2 with an alkyl halide (such as benzyl bromide or 4-chlorobenzyl chloride) and potassium carbonate in DMF gives 11-Scheme 2. Triazole 11-Scheme 2 is protected as the methoxy methylethyl ether to afford 12-Scheme 2. Alkylation of 12-Scheme 2 with an alkyl halide (such as methyl iodide, ethyl iodide, *i*-isobutyl iodide, *n*-propyl iodide, butyl iodide, allyl bromide, benzyl bromide, and methyl bromoacetate) affords the desired tertiary amine 13-Scheme 2. Deprotection of the MOM-ether 13-Scheme 2 with trifluoroacetic acid (TFA) provides the desired product 14-Scheme 2.

Formulation of Pharmaceutical Compositions

The pharmaceutically effective compounds of this invention (and the pharmaceutically acceptable salts thereof) are administered in conventional dosage forms prepared by combining a compound of this invention ("active ingredient") in an amount sufficient to treat cancer, haemangioma, proliferative retinopathy, rheumatoid arthritis, atherosclerotic

neovascularization, psoriasis, ocular neovascularization or obesity ("MetAp2-mediated disease states") with standard pharmaceutical carriers or diluents according to conventional procedures well known in the art. These procedures may involve mixing, granulating and compressing or dissolving the

5 ingredients as appropriate to the desired preparation.

The pharmaceutical carrier employed may be, for example, either a solid or liquid. Exemplary of solid carriers are lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid and the like. Exemplary of liquid carriers are syrup, peanut oil, olive oil, water and the like.

10 Similarly, the carrier or diluent may include time delay material well known to the art, such as glyceryl monostearate or glyceryl distearate alone or with a wax.

A wide variety of pharmaceutical forms can be employed. Thus, if a solid carrier is used, the preparation can be tableted, placed in a hard gelatin capsule in powder or pellet form or in the form of a troche or lozenge. The amount of solid carrier will vary widely but preferably will be from about 25 mg to about 1000 mg. When a liquid carrier is used, the preparation will be in the form of a syrup, emulsion, soft gelatin capsule, sterile injectable liquid such as an ampule or nonaqueous liquid suspension.

15 20 The active ingredient may also be administered topically to a mammal in need of treatment or prophylaxis of MetAP2-mediated disease states. The amount of active ingredient required for therapeutic effect on topical administration will, of course, vary with the compound chosen, the nature and severity of the disease state being treated and the mammal undergoing treatment, and is ultimately at the discretion of the physician. A suitable dose of an active ingredient is 1.5 mg to 500 mg for topical administration, the most preferred dosage being 1 mg to 100 mg, for example 5 to 25 mg administered two or three times daily.

By topical administration is meant non-systemic administration and includes the application of the active ingredient externally to the epidermis, to the buccal cavity and instillation of such a compound into the ear, eye and nose, and where the compound does not significantly enter the blood stream. By systemic administration is meant oral, intravenous, intraperitoneal and intramuscular administration.

35 While it is possible for an active ingredient to be administered alone as the raw chemical, it is preferable to present it as a pharmaceutical formulation. The active ingredient may comprise, for topical administration, from 0.001% to

10% w/w, e.g. from 1% to 2% by weight of the formulation although it may comprise as much as 10% w/w but preferably not in excess of 5% w/w and more preferably from 0.1% to 1% w/w of the formulation.

The topical formulations of the present invention, both for veterinary and
5 for human medical use, comprise an active ingredient together with one or more acceptable carrier(s) therefor and optionally any other therapeutic ingredient(s). The carrier(s) must be 'acceptable' in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

Formulations suitable for topical administration include liquid or semi-
10 liquid preparations suitable for penetration through the skin to the site of inflammation such as liniments, lotions, creams, ointments or pastes, and drops suitable for administration to the eye, ear or nose.

Drops according to the present invention may comprise sterile aqueous or oily solutions or suspensions and may be prepared by dissolving the active
15 ingredient in a suitable aqueous or alcoholic solution of a bactericidal and/or fungicidal agent and/or any other suitable preservative, and preferably including a surface active agent. The resulting solution may then be clarified by filtration, transferred to a suitable container which is then sealed and sterilized by autoclaving or maintaining at 98-100°C for half an hour. Alternatively, the
20 solution may be sterilized by filtration and transferred to the container by an aseptic technique. Examples of bactericidal and fungicidal agents suitable for inclusion in the drops are phenylmercuric nitrate or acetate (0.002%), benzalkonium chloride (0.01%) and chlorhexidine acetate (0.01%). Suitable solvents for the preparation of an oily solution include glycerol, diluted alcohol
25 and propylene glycol.

Lotions according to the present invention include those suitable for application to the skin or eye. An eye lotion may comprise a sterile aqueous solution optionally containing a bactericide and may be prepared by methods
30 similar to those for the preparation of drops. Lotions or liniments for application to the skin may also include an agent to hasten drying and to cool the skin, such as an alcohol or acetone, and/or a moisturizer such as glycerol or an oil such as castor oil or arachis oil.

Creams, ointments or pastes according to the present invention are semi-solid formulations of the active ingredient for external application. They may be
35 made by mixing the active ingredient in finely divided or powdered form, alone or in solution or suspension in an aqueous or non-aqueous fluid, with the aid of suitable machinery, with a greasy or non-greasy basis. The basis may comprise

hydrocarbons such as hard; soft or liquid paraffin, glycerol, beeswax, a metallic soap; a mucilage; an oil of natural origin such as almond, corn, arachis, castor or olive oil; wool fat or its derivatives, or a fatty acid such as stearic or oleic acid together with an alcohol such as propylene glycol. The formulation may
5 incorporate any suitable surface-active agent such as an anionic, cationic or non-ionic surfactant such as esters or polyoxyethylene derivatives thereof. Suspending agents such as natural gums, cellulose derivatives or inorganic materials such as siliceous silicas, and other ingredients such as lanolin, may also be included.

10 The active ingredient may also be administered by inhalation. By "inhalation" is meant intranasal and oral inhalation administration. Appropriate dosage forms for such administration, such as an aerosol formulation or a metered dose inhaler, may be prepared by conventional techniques. The daily dosage amount of the active ingredient administered by inhalation is from about
15 0.1 mg to about 100 mg per day, preferably about 1 mg to about 10 mg per day.

In one aspect, this invention relates to a method of treating cancer, haemangioma, proliferative retinopathy, rheumatoid arthritis, atherosclerotic neovascularization, psoriasis, ocular neovascularization or obesity, all in mammals, preferably humans, which comprises administering to such
20 mammal an effective amount of a MetAP2 inhibitor, in particular, a compound of this invention.

By the term "treating" is meant either prophylactic or therapeutic therapy. Such compound can be administered to such mammal in a conventional dosage form prepared by combining the compound of this
25 invention with a conventional pharmaceutically acceptable carrier or diluent according to known techniques. It will be recognized by one of skill in the art that the form and character of the pharmaceutically acceptable carrier or diluent is dictated by the amount of active ingredient with which it is to be combined, the route of administration and other well-known variables. The
30 compound is administered to a mammal in need of treatment for cancer, haemangioma, proliferative retinopathy, rheumatoid arthritis, atherosclerotic neovascularization, psoriasis, ocular neovascularization or obesity, in an amount sufficient to decrease symptoms associated with these disease states. The route of administration may be oral or parenteral.

35 The term parenteral as used herein includes intravenous, intramuscular, subcutaneous, intra-rectal, intravaginal or intraperitoneal administration. The subcutaneous and intramuscular forms of parenteral administration are

generally preferred. The daily parenteral dosage regimen will preferably be from about 30 mg to about 300 mg per day of active ingredient. The daily oral dosage regimen will preferably be from about 100 mg to about 2000 mg per day of active ingredient.

5 It will be recognized by one of skill in the art that the optimal quantity and spacing of individual dosages of a compound of this invention will be determined by the nature and extent of the condition being treated, the form, route and site of administration, and the particular mammal being treated, and that such optimums can be determined by conventional techniques. It will
10 also be appreciated by one of skill in the art that the optimal course of treatment, i.e., the number of doses of the compound given per day for a defined number of days, can be ascertained by those skilled in the art using conventional course of treatment determination tests.

15 EXAMPLES

The invention will now be described by reference to the following examples which are merely illustrative and are not to be construed as a limitation of the scope of the present invention. In the Examples, proton NMR spectra were performed upon a Bruker 400 MHz NMR spectrometer, unless
20 otherwise indicated.

Example 1

Preparation of 3-anilino-5-benzylthio-1,2,4-triazole

a) 1-Phenyl-2,4-dithiobiuret

To a stirring solution of NaOH (0.52 g, 13.1 mmol) in 60 mL of 10%
25 H₂O:CH₃CN was added thiourea (1.0 g, 13.1 mmol). The resulting suspension was heated to 40 °C for 20 min. and then cooled to RT. To this mixture was added phenylisothiocyanate (1.5 ml, 13.1 mmol), and the clear yellow solution was stirred overnight. After stirring for 12 h, 1 N HCl was added until a white precipitate formed. The precipitate was filtered, washed with H₂O, and dried
30 under vacuum to produce the title compound as a light yellow powder (0.78 g, 30%). ¹H-NMR (400MHz, d6-DMSO) δ7.25 (t, 2H, J=7.3 Hz), 7.40 (t, 2H, J=7.9 Hz), 7.56 (d, 1H, J=7.9 Hz), 9.13-9.29 (broad singlet, 1H), and 10.26-10.79 (broad singlet, 2H).

b) 2-Ethyl-1-phenyl-2-isodithiobiuret

35 To a stirring solution of the compound of Example 1(a) (150 mg, 0.70 mmol) in 4 mL of DMF was added triethylamine (57 uL, 0.70 mmol). The resulting yellow/green solution was stirred for 10 min at RT. To this solution

was added ethyl iodide (100 uL, 0.70 mmol), and the reaction mixture was stirred for 2 h at RT. The yellow solution was poured into 20 mL of H₂O and extracted four times with EtOAc. The organic extracts were dried over Na₂SO₄, filtered, concentrated, and the crude residue was subjected to column chromatography (silica gel; ethyl acetate/hexane) to afford the title compound as a white crystalline solid (108 mg, 64%). ¹H-NMR (400MHz, d₆-DMSO) δ 1.22 (t, 3H, J=7.2 Hz), 2.96 (quartet, 2H, J=7.2 Hz), 6.85 (d, 1H, J=7.6 Hz), 7.16 (t, 1H, J=7.2 Hz), 7.29-7.41 (m, 3H), 8.27 (broad singlet, 1H), 9.89 (broad singlet, 1H), and 10.99 (broad singlet, 1H).

c) 3-anilino-5-mercapto-1,2,4-triazole

To a stirring solution of the compound of Example 1(b) in 2 mL of EtOH was added 50 uL of anhydrous hydrazine. The reaction mixture was heated at 80 °C for 1 h, concentrated to dryness, and then purified by preparative HPLC to yield the title compound as a white solid (30 mg, 37%). MS (ESI) 190.90 (M-H)⁺.

d) 3-anilino-5-benzylthio-1,2,4-triazole

To a stirring solution of the compound of Example 1(c) (23 mg, 0.12 mmol) in 1.2 mL of DMF was added K₂CO₃ (17 mg, 0.12 mmol), followed by benzyl bromide (20 mg, 0.12 mmol). The mixture was stirred overnight, filtered, and purified by preparative HPLC to afford the title compound as a white solid (30 mg, 70%). ¹H-NMR (400MHz, d₆-DMSO) δ 9.30 (broad singlet, 1H), 7.47 (d, 2H, J=8.1 Hz), 7.39 (d, 2H, J=7.3 Hz), 7.31 (t, 2H, J=7.3 Hz), 7.23 (quartet, 3H, J=7.3 Hz), 6.82 (t, 1H, J=7.3 Hz), and 4.3 (s, 2H).

Example 2

Preparation of 3-anilino-5-(4-chlorobenzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except substituting 4-chlorobenzyl bromide for benzyl bromide in step 1(d), the title compound was prepared as a white solid. ¹H-NMR (400MHz, d₆-DMSO) δ 9.32 (broad singlet, 1H), 7.46 (d, 2H, J=7.8 Hz), 7.41 (d, 2H, J=8.4 Hz), 7.36 (d, 2H, J=8.4 Hz), 7.22 (t, 2H, J=7.8 Hz), 6.82 (t, 1H, J=7.24 Hz), and 4.33 (s, 2H).

Example 3

Preparation of 3-anilino-5-methylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except methyl iodide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 413.2 (2M+H)⁺.

Example 4

Preparation of 3-anilino-5-allylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except allyl bromide
5 was substituted for benzyl bromide in step 1(d), the title compound was
prepared as a white solid. MS (ESI) 233.0 (M+H)⁺.

Example 5

Preparation of 3-anilino-5-(2-methyl-2-butenylthio)-1,2,4-triazole

10 Following the procedure of Example 1(a)-1(d), except 1-bromo-3-
methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title
compound was prepared as a white solid. MS (ESI) 261.2 (M+H)⁺.

Example 6

Preparation of 3-anilino-5-(2-methyl-butylthio)-1,2,4-triazole

15 Following the procedure of Example 1(a)-1(d), except 1-bromo-3-
methylbutane was substituted for benzyl bromide in step 1(d), the title
compound was prepared as a white solid. MS (ESI) 263.2 (M+H)⁺.

Example 7

Preparation of 3-anilino-5-(2-methyl-2-pentenylthio)-1,2,4-triazole

20 Following the procedure of Example 1(a)-1(d), except 5-bromo-2-
methyl-2-pentene was substituted for benzyl bromide in step 1(d), the title
compound was prepared as a white solid. MS (ESI) 275.2 (M+H)⁺.

25

Example 8

Preparation of 3-anilino-5-(α -methylbenzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except (1-bromoethyl)
benzene was substituted for benzyl bromide in step 1(d), the title compound
30 was prepared as a white solid. MS (ESI) 297.2 (M+H)⁺.

Example 9

Preparation of 3-anilino-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except
35 bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the
title compound was prepared as a white solid. MS (ESI) 289.0 (M+H)⁺.

Example 10

Preparation of 3-anilino-5-(propyl acetylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except propyl bromoacetate was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 293.2 (M+H)⁺.

Example 11

Preparation of 3-anilino-5-(3,3-dimethoxy-propylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-bromo-1,1-dimethoxy-propane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 295.2 (M+H)⁺.

Example 12

Preparation of 3-anilino-5-(2-phenethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except (2-bromoethyl)benzene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 297.2 (M+H)⁺.

Example 13

Preparation of 3-anilino-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 288.2 (M+H)⁺.

Example 14

Preparation of 3-anilino-5-(3-phenyl-[1,2,4]oxadiazol-5-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-chloromethyl-5-phenyl-1,2,4-oxadiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 351.2 (M+H)⁺.

Example 15

Preparation of 3-anilino-5-(1H-benzimidazol-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-(chloromethyl)-benzimidazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 323.2 (M+H)⁺.

Example 16Preparation of 3-anilino-5-(2-(4-chlorophenyl)-thiazol-4-ylmethylthio)-1,2,4-triazole

5 Following the procedure of Example 1(a)-1(d), except 4-chloromethyl-2-(4-chlorophenyl)thiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 400.0 (M+H)⁺.

Example 17

10 Preparation of 3-anilino-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-chloromethyl-2-methylthiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 304.2 (M+H)⁺.

Example 18

15 Preparation of 3-anilino-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 284.2 (M+H)⁺.

Example 19

20 Preparation of 3-anilino-5-(pyridin-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 284.0 (M+H)⁺.

Example 20

30 Preparation of 3-anilino-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 289.0 (M+H)⁺.

Example 21

35 Preparation of 3-anilino-5-(4-*i*-propyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-isopropylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 325.2 (M+H)⁺.

Example 22Preparation of 3-anilino-5-(quinolin-8-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 8-bromomethylquinoline was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 334.2 (M+H)⁺.

Example 23Preparation of 3-anilino-5-(4-acetamido-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-acetamidobenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 340.2 (M+H)⁺.

Example 24Preparation of 4-(5-anilino-2 H-[1,2,4]triazol-3-yl thio)-benzoic acid

Following the procedure of Example 1(a)-1(d), except 4-(chloromethyl)benzoic acid was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 327.2 (M+H)⁺.

Example 25Preparation of 3-anilino-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 297.0 (M+H)⁺.

Example 26Preparation of 3-anilino-5-(4-trifluoromethyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-(trifluoromethyl)benzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 350.8 (M+H)⁺.

Example 27Preparation of 3-anilino-5-(3,5-dimethyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,5-dimethylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 311.4 (M+H)⁺.

Example 28

Preparation of 3-anilino-5-(4-cyano-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-cyanobenzyl
5 bromide was substituted for benzyl bromide in step 1(d), the title compound
was prepared as a white solid. MS (ESI) 308.2 (M+H)⁺.

Example 29

Preparation of 3-anilino-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

10 Following the procedure of Example 1(a)-1(d), except 3,4-
difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title
compound was prepared as a white solid. MS (ESI) 319.0 (M+H)⁺.

Example 30

Preparation of 3-anilino-5-(furan-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-chloromethyl-
furan (Berry, J. M.; Watson, C. Y.; Whish, W. J. D.; Threadgill, M. D. J.
Chem. Soc. Perkin Trans. 1 **1997**, 8, 1147) was substituted for benzyl bromide
in step 1(d), the title compound was prepared as a white solid. MS (ESI)
20 273.2 (M+H)⁺.

Example 31

Preparation of 3-anilino-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-chloromethyl-
25 3-methyl-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R.
C. Tetrahedron Lett. **1988**, 29(1), 117) was substituted for benzyl bromide in
step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2
(M+H)⁺.

Example 32

Preparation of 3-anilino-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-chloromethyl-
3-chloro-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R.
C. Tetrahedron Lett. **1988**, 29(1), 117) was substituted for benzyl bromide in
35 step 1(d), the title compound was prepared as a white solid. MS (ESI) 323.2
(M+H)⁺.

Example 33Preparation of 3-anilino-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-chloromethyl-5-methyl-thiophene (Moradpour, A. *J. Chem. Soc. Perkin Trans. 1*, **1993**, 1, 7) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 34Preparation of 3-anilino-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-chloromethyl-5-chloro-thiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 323.0 (M+H)⁺.

Example 35Preparation of 5-(5-phenylamino-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester

Following the procedure of Example 1(a)-1(d), except 5-chloromethyl-furan-2-carboxylic acid ethyl ester was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 345.2 (M)⁺.

Example 36Preparation of 3-anilino-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-chloromethyl-5-bromo-thiophene (Clapp, R. C.; Clark, J. H.; Vaughan, J. R.; English, J. P.; Anderson, G. W. *J. Am. Chem. Soc.* **1947**, 60, 1549) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 367.0 (M)⁺.

Example 37Preparation of 5-(5-phenylamino-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde

Following the procedure of Example 1(a)-1(d), except 5-chloromethyl-furan-2-carbaldehyde (Sanda, K.; Rigal, L.; Delmas, M.; Gaset, A. *Synthesis* **1992**, 6, 541) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 301.2 (M+H)⁺.

Example 38

Preparation of 3-anilino-5-(thiophen-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-chloromethylthiophene (Lamy, J.; Lavit, D.; Buu-Hoi, N. P. *J. Chem. Soc.* **1958**, 4202) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 289.0 (M+H)⁺.

Example 39

Preparation of 3-anilino-5-(furan-3-ylthio)-1,2,4-triazole ..

Following the procedure of Example 1(a)-1(d), except 3-chloromethylfuran (Arena, G.; Cali, R.; Maccarone, E.; Passerini, A. *J. Chem. Soc. Perkin Trans. 2* **1993**, 10, 1941) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 273.2 (M+H)⁺.

Example 40

Preparation of 3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a), the title compound was prepared as a white solid. MS (ESI) 297.0 (M+H)⁺.

Example 41

Preparation of 3-(4-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 42

Preparation of 3-(4-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.0 (M+H)⁺.

Example 43Preparation of 3-(4-methyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 298.2 (M+H)⁺.

Example 44Preparation of 3-(4-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 275.2 (M+H)⁺.

Example 45Preparation of 3-(4-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.0 (M+H)⁺.

Example 46Preparation of 3-(4-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 302.2 (M+H)⁺.

Example 47Preparation of 3-(4-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 311.2 (M+H)⁺.

Example 48

Preparation of 3-(4-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 333.2 (M+H)⁺.

Example 49

Preparation of 3-(4-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 327.2 (M+H)⁺.

Example 50

Preparation of 3-(4-methyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-chloromethyl-2-methylthiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 318.2 (M+H)⁺.

Example 51

Preparation of 3-(4-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 298.2 (M+H)⁺.

Example 52

Preparation of 3-(4-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-furan (Berry, J. M.; Watson, C. Y.; Whish, W. J. D.; Threadgill, M. D. *J. Chem. Soc. Perkin Trans. 1* 1997, 8, 1147) was substituted for benzyl

bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 287.2 (M+H)⁺.

Example 53

5 Preparation of 3-(4-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-methyl-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* 1988, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 317.2 (M+H)⁺.

Example 54

15 Preparation of 3-(4-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-chloro-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* 1988, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 337.2 (M+H)⁺.

Example 55

25 Preparation of 3-(4-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-methyl-thiophene (Moradpour, A. J. *Chem. Soc. Perkin Trans. 1*, 1993, 1, 7) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 317.2 (M+H)⁺.

Example 56

35 Preparation of 3-(4-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-

chloromethyl-5-chloro-thiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 337.2 (M+H)⁺.

5

Example 57Preparation of 5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carboxylic acid ethyl ester was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 359.2 (M+H)⁺.

10

Example 58Preparation of 3-(4-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-bromo-thiophene (Clapp, R. C.; Clark, J. H; Vaughan, J. R.; English, J. P.; Anderson, G. W. *J. Am. Chem. Soc.* **1947**, *60*, 1549) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 381.0 (M)⁺.

20

Example 59Preparation of 5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carbaldehyde (Sanda, K.; Rigal, L.; Delmas, M.; Gaset, A. *Synthesis* **1992**, *6*, 541) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.2 (M+H)⁺.

30

Example 60Preparation of 3-(4-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-thiophene (Lamy, J.; Lavit, D.; Buu-Hoi, N. P. *J. Chem. Soc.*

35

1958, 4202) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 61

5 Preparation of 3-(4-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-furan (Arena, G.; Cali, R.; Maccarone, E.; Passerini, A. *J. Chem. Soc. Perkin Trans. 2* 1993, 10, 1941) was substituted for benzyl bromide in
10 step 1(d), the title compound was prepared as a white solid. MS (ESI) 287.2 (M+H)⁺.

Example 62

15 Preparation of 3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 297.2 (M+H)⁺.

Example 63

20 Preparation of 3-(2-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1 (d), the
25 title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 64

30 Preparation of 3-(2-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 65

35 Preparation of 3-(2-methyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d) except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-

(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 298.2 (M+H)⁺.

Example 66

5 Preparation of 3-(2-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 275.2 (M+H)⁺.

10

Example 67

Preparation of 3-(2-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.2 (M+H)⁺.

15

Example 68

20 Preparation of 3-(2-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 302.0 (M+H)⁺.

25

Example 69

Preparation of 3-(2-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 311.2 (M+H)⁺.

30

Example 70

35 Preparation of 3-(2-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-

difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 333.2 (M+H)⁺.

Example 71

5 Preparation of 3-(2-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 327.2 (M+H)⁺.

10

Example 72

Preparation of 3-(2-methyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-chloromethyl-2-methylthiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 318.2 (M+H)⁺.

20

Example 73

Preparation of 3-(2-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 298.2 (M+H)⁺.

25

Example 74

Preparation of 3-(2-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-furan (Berry, J. M.; Watson, C. Y.; Whish, W. J. D.; Threadgill, M. D. *J. Chem. Soc. Perkin Trans. 1* **1997**, 8, 1147) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 287.2 (M+H)⁺.

35

Example 75Preparation of 3-(2-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-methyl-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 317.2 (M+H)⁺.

10

Example 76Preparation of 3-(2-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-chloro-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 337.2 (M+H)⁺.

20

Example 77Preparation of 3-(2-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-methyl-thiophene (Moradpour, A. J. *Chem. Soc. Perkin Trans. 1*, **1993**, 1, 7) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 317.2 (M+H)⁺.

30

Example 78Preparation of 3-(2-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-chloro-thiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 337.2 (M+H)⁺.

35

Example 79

Preparation of 5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester

5 Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carboxylic acid ethyl ester was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 359.2 (M+H)⁺.

10

Example 80

Preparation of 3-(2-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole

15 Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-bromo-thiophene (Clapp, R. C.; Clark, J. H; Vaughan, J. R.; English, J. P.; Anderson, G. W. *J. Am. Chem. Soc.* **1947**, *60*, 1549) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid . MS (ESI) 381.0 (M)⁺.

20

Example 81

Preparation of 5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde

25 Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carbaldehyde (Sanda, K.; Rigal, L.; Delmas, M.; Gaset, A. *Synthesis* **1992**, *6*, 541) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.2 (M+H)⁺.

30

Example 82

Preparation of 3-(2-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole

35 Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-thiophene (Lamy, J.; Lavit, D.; Buu-Hoi, N. P. *J. Chem. Soc.* **1958**, 4202) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 83Preparation of 3-(2-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-furan (Arena, G.; Cali, R.; Maccarone, E.; Passerini, A. *J. Chem. Soc. Perkin Trans. 2* **1993**, *10*, 1941) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 287.2 (M+H)⁺.

Example 84Preparation of 3-(4-chloro-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a), the title compound was prepared as a white solid. MS (ESI) 317.2 (M+H)⁺.

Example 85Preparation of 3-(4-chloro-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 322.7 (M)⁺.

Example 86Preparation of 3-(4-chloro-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 323.2 (M+H)⁺.

Example 87Preparation of 3-(4-chloro-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 318.2 (M+H)⁺.

Example 88

Preparation of 3-(4-chloro-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 295.2 (M+H)⁺.

Example 89

Preparation of 3-(4-chloro-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 335.2 (M+H)⁺.

Example 90

Preparation of 3-(4-chloro-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 322.0 (M+H)⁺.

Example 91

Preparation of 3-(4-chloro-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 331.0 (M+H)⁺.

Example 92

Preparation of 3-(4-chloro-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 352.8 (M)⁺.

Example 93Preparation of 3-(4-chloro-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 347.0 (M+H)⁺.

Example 94Preparation of 3-(4-chloro-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-chloromethyl-2-methylthiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 338.2 (M+H)⁺.

Example 95Preparation of 3-(4-chloro-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 318.0 (M+H)⁺.

Example 96Preparation of 3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a), the title compound was prepared as a white solid. MS (ESI) 313.2 (M+H)⁺.

Example 97Preparation of 3-(4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 319.0 (M+H)⁺.

Example 98Preparation of 3-(4-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 319.2 (M+H)⁺.

Example 99Preparation of 3-(4-methoxy-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 314.2 (M+H)⁺.

Example 100Preparation of 3-(4-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 291.2 (M+H)⁺.

Example 101Preparation of 3-(4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 331.2 (M+H)⁺.

Example 102Preparation of 3-(4-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 318.2 (M+H)⁺.

Example 103

Preparation of 3-(4-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 327.2 (M+H)⁺.

Example 104

Preparation of 3-(4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 349.0 (M+H)⁺.

Example 105

Preparation of 3-(4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 343.0 (M+H)⁺.

Example 106

Preparation of 3-(4-methoxy-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-chloromethyl-2-methylthiazole was substituted for benzyl

bromide in step 1(d), the title compound was prepared as a white solid (2%).
MS (ESI) 334.2 (M+H)⁺.

Example 107

5 Preparation of 3-(4-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 314.2
10 (M+H)⁺.

Example 108

Preparation of 3-(4-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole

15 Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-chloro-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared
20 as a white solid . MS (ESI) 353.2 (M+H)⁺.

Example 109

Preparation of 3-(4-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole

25 Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-chloro-thiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid . MS (ESI) 353.0 (M+H)⁺.

30

Example 110

Preparation of 4-(5-benzylthio-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

35 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a), the title compound was prepared as a white solid. MS (ESI) 341.0 (M+H)⁺.

Example 111Preparation of 4-(5-(cyclohexylmethylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

5 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 347.2 (M+H)⁺.

10

Example 112Preparation of 4-(5-(pyridin-4-ylmethylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

15 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 342.2 (M+H)⁺.

20

Example 113Preparation of 4-(5-(2-methyl-2-butenylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

25 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 319.0 (M+H)⁺.

Example 114

30 Preparation of 4-(5-(2-fluoro-benzylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

35 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 359.2 (M+H)⁺.

Example 115Preparation of 4-(5-(5-methyl-isoxazol-3-ylmethylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

5 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 346.0 (M+H)⁺.

10

Example 116Preparation of 4-(5-(2-methyl-benzylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

15 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 355.0 (M+H)⁺.

20

Example 117Preparation of 4-(5-(3-methoxy-benzylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

25 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 371.0 (M+H)⁺.

Example 118

30 Preparation of 4-(5-(3,4-difluoro-benzylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

35 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 377.0 (M+H)⁺.

Example 119Preparation of 4-(5-(2-methoxy-benzylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 371.0 (M+H)⁺.

Example 120Preparation of 4-(5-(2-methyl-thiazol-4-ylmethylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-chloromethyl-2-methylthiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 362.0 (M+H)⁺.

Example 121Preparation of 4-(5-(pyridin-2-ylmethylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 342.2 (M+H)⁺.

Example 122Preparation of 3-(3,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a), the title compound was prepared as a white solid. MS (ESI) 343.0 (M+H)⁺.

Example 123Preparation of 3-(3,4-dimethoxy-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-methoxybenzyl chloride was substituted for benzyl bromide in step 1 (d), the title compound was prepared as a white solid. MS (ESI) 373.2 (M+H)⁺.

Example 124

Preparation of 3-(3,4-dimethoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 349.2 (M+H)⁺.

Example 125

Preparation of 3-(3,4-dimethoxy-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 344.2 (M+H)⁺.

Example 126

Preparation of 3-(3,4-dimethoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 321.2 (M+H)⁺.

Example 127

Preparation of 3-(3,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 361.2 (M+H)⁺.

Example 128

Preparation of 3-(3,4-dimethoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 348.2 (M+H)⁺.

Example 129

Preparation of 3-(3,4-dimethoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 357.2 (M+H)⁺.

Example 130

Preparation of 3-(3,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 379.0 (M+H)⁺.

Example 131

Preparation of 3-(3,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 373.0 (M+H)⁺.

Example 132

Preparation of 3-(3,4-dimethoxy-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-chloromethyl-2-methylthiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 364.2 (M+H)⁺.

Example 133

Preparation of 3-(3,4-dimethoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 344.0 (M+H)⁺.

Example 134

Preparation of 3-(3,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 349.0 (M+H)⁺.

Example 135

Preparation of 3-(2-phenyl-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 359.2 (M+H)⁺.

Example 136Preparation of 3-(2-phenyl-anilino)-5-(3-methoxybenzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl
5 isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-methoxyphenyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 389.0 (M+H)⁺.

Example 137

10 Preparation of 3-(2-phenyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl
isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and
bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the
title compound was prepared as a white solid. MS (ESI) 365.2 (M+H)⁺.

15

Example 138Preparation of 3-(2-phenyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d) except 2-phenyl-phenyl
isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-
20 (chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 360.2 (M+H)⁺.

Example 139Preparation of 3-(2-phenyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

25 Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 337.2 (M+H)⁺.

30

Example 140Preparation of 3-(2-phenyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl
isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title
35 compound was prepared as a white solid. MS (ESI) 376.8 (M)⁺.

Example 141Preparation of 3-(2-phenyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl
5 isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 364.0 (M+H)⁺.

Example 142Preparation of 3-(2-phenyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl
isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the
15 title compound was prepared as a white solid. MS (ESI) 373.0 (M+H)⁺.

Example 143Preparation of 3-(2-phenyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl
20 isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 395.0 (M+H)⁺.

Example 144Preparation of 3-(2-phenyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl
isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl bromide was substituted for benzyl bromide in step 1(d), the
title compound was prepared as a white solid. MS (ESI) 389.2 (M+H)⁺.

30

Example 145Preparation of 3-(2-phenyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl
35 isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-chloromethyl-2-methylthiazole was substituted for benzyl bromide in step

1(d), the title compound was prepared as a white solid. MS (ESI) 380.0 (M+H)⁺.

Example 146

5 Preparation of 3-(2-phenyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 365.2 (M+H)⁺.

10

Example 147

Preparation of [5-(benzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 284.2 (M+H)⁺.

15

Example 148

Preparation of [5-(3-methoxybenzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-methoxyphenyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 314.2 (M+H)⁺.

20

Example 149

Preparation of [5-(cyclohexylmethylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 290.2 (M+H)⁺.

30

Example 150

Preparation of [5-(pyridin-4-ylmethylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

35

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-

(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 285.2 (M+H)⁺.

Example 151

5 Preparation of [5-(2-methyl-2-butenylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the
10 title compound was prepared as a white solid. MS (ESI) 262.0 (M+H)⁺.

Example 152

Preparation of [5-(2-fluoro-benzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

15 Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 302.2 (M+H)⁺.

Example 153

20 Preparation of [5-(5-methyl-isoxazol-3-yl)methylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step
25 1(d), the title compound was prepared as a white solid. MS (ESI) 289.0 (M+H)⁺.

Example 154

30 Preparation of [5-(2-methyl-benzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the
35 title compound was prepared as a white solid. MS (ESI) 298.2 (M+H)⁺.

Example 155Preparation of [5-(3,4-difluoro-benzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 320.2 (M+H)⁺.

Example 156

10 Preparation of [5-(2-methoxy-benzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 314.2 (M+H)⁺.

Example 157

20 Preparation of [5-(pyridin-2-ylmethylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 285.2 (M+H)⁺.

Example 158

25 Preparation of [5-(2-methyl-thiazol-4-ylmethylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-chloromethyl-2-methylthiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 305.2 (M+H)⁺.

Example 159

35 Preparation of [5-(thiophen-2-ylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 290.0 (M+H)⁺.

5

Example 160

Preparation of 3-(2-ethyl-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 311.4 (M+H)⁺.

10

Example 161

Preparation of 3-(2-ethyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1 (d), the title compound was prepared as a white solid. MS (ESI) 317.2 (M+H)⁺.

15

Example 162

Preparation of 3-(2-ethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 329.2 (M+H)⁺.

20

Example 163

Preparation of 3-(2-ethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 347.0 (M+H)⁺.

30

Example 164

Preparation of 3-(2-ethyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-

35

bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 289.0 (M+H)⁺.

Example 165

5 Preparation of 3-(2-ethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 329.2 (M+H)⁺.

10

Example 166

Preparation of 3-(2-ethyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 325.2 (M+H)⁺.

15

Example 167

20 Preparation of 3-(2-ethyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chlorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 345.0 (M+H)⁺.

25

Example 168

Preparation of 3-(2-ethyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 341.2 (M+H)⁺.

30

Example 169

35 Preparation of 3-(2-ethyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-

methylenedioxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 355.0 (M+H)⁺.

5

Example 170

Preparation of 3-(2-ethyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 316.0 (M+H)⁺.

15

Example 171

Preparation of 3-(2-ethyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 312.2 (M+H)⁺.

20

Example 172

Preparation of 3-(2-ethyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 341.2 (M+H)⁺.

25

Example 173

Preparation of 3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 313.2 (M+H)⁺.

35

Example 174

Preparation of 3-(2-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 319.0 (M+H)⁺.

Example 175

Preparation of 3-(2-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 331.2 (M+H)⁺.

Example 176

Preparation of 3-(2-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 319.2 (M+H)⁺.

Example 177

Preparation of 3-(2-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 349.0 (M+H)⁺.

Example 178

Preparation of 3-(2-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide

in step 1(d), the title compound was prepared as a white solid. MS (ESI) 291.2 (M+H)⁺.

Example 179

5 Preparation of 3-(2-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 331.2 (M+H)⁺.

Example 180

15 Preparation of 3-(2-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 327.2 (M+H)⁺.

Example 181

20 Preparation of 3-(2-methoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chlorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 347.0 (M+H)⁺.

Example 182

30 Preparation of 3-(2-methoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 343.0 (M+H)⁺.

35

Example 183Preparation of 3-(2-methoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-methylenedioxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 357.0 (M+H)⁺.

Example 184Preparation of 3-(2-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 316.0 (M+H)⁺.

Example 185Preparation of 3-(2-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 314.2 (M+H)⁺.

Example 186Preparation of 3-(2-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid (49%). MS (ESI) 343.0 (M+H)⁺.

Example 187Preparation of 3-(2-methoxy-anilino)-5-(furan-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-furan (Berry, J. M.; Watson, C. Y.; Whish, W. J. D.; Threadgill, M. D. *J. Chem. Soc. Perkin Trans. 1* **1997**, 8, 1147) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 188

Preparation of 3-(2-methoxy-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-methyl-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 333.2 (M+H)⁺.

Example 189

Preparation of 3-(2-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-chloro-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 353.2 (M+H)⁺.

Example 190

Preparation of 3-(2-methoxy-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-methyl-thiophene (Moradpour, A. *J. Chem. Soc. Perkin Trans. 1*, **1993**, 1, 7) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 333.2 (M+H)⁺.

Example 191Preparation of 3-(2-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-chloro-thiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 353.0 (M+H)⁺.

Example 192Preparation of 5-(5-(2-methoxyphenylamino)-4H-[1,2,4]triazol-3-yl)sulfanylmethyl)-furan-2-carboxylic acid ethyl ester

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carboxylic acid ethyl ester was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 375.0 (M+H)⁺.

Example 193Preparation of 3-(2-methoxy-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-bromo-thiophene (Clapp, R. C.; Clark, J. H.; Vaughan, J. R.; English, J. P.; Anderson, G. W. *J. Am. Chem. Soc.* **1947**, 60, 1549) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 396.8 (M-H)⁺.

Example 194Preparation of 3-(2-methoxy-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-thiophene (Lamy, J.; Lavit, D.; Buu-Hoi, N. P. *J. Chem. Soc.* **1958**, 4202) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 319.0 (M+H)⁺.

Example 195Preparation of 3-(2-methoxy-anilino)-5-(furan-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in
5 step 1(a) and 3-chloromethyl-furan (Arena, G.; Cali, R.; Maccarone, E.; Passerini, A. *J. Chem. Soc. Perkin Trans. 2* **1993**, *10*, 1941) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 196Preparation of 3-(2-isopropyl-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in
10 step 1(a) the title compound was prepared as a white solid. MS (ESI) 325.2 (M+H)⁺.

Example 197Preparation of 3-(2-isopropyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in
20 step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 331.2 (M+H)⁺.

Example 198Preparation of 3-(2-isopropyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in
25 step 1(a) and 4-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 343.0 (M+H)⁺.

Example 199Preparation of 3-(2-isopropyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in
35 step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in

step 1(d), the title compound was prepared as a white solid. MS (ESI) 331.2 (M+H)⁺.

Example 200

5 Preparation of 3-(2-isopropyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide
10 in step 1(d), the title compound was prepared as a white solid. MS (ESI) 361.2 (M+H)⁺.

Example 201

15 Preparation of 3-(2-isopropyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide
20 in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 202

Preparation of 3-(2-isopropyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in
25 step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 343.0 (M+H)⁺.

Example 203

30 Preparation of 3-(2-isopropyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in
35 step 1(d), the title compound was prepared as a white solid. MS (ESI) 339.2 (M+H)⁺.

Example 204Preparation of 3-(2-isopropyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chlorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 359.2 (M+H)⁺.

Example 205Preparation of 3-(2-isopropyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 355.2 (M+H)⁺.

Example 206Preparation of 3-(2-isopropyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-methylenedioxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 369.2 (M+H)⁺.

Example 207Preparation of 3-(2-isopropyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylisothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 330.2 (M+H)⁺.

Example 208Preparation of 3-(2-isopropyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 326.2 (M+H)⁺.

Example 209Preparation of 3-(2-isopropyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 355.2 (M+H)⁺.

Example 210Preparation of 3-(2-isopropyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-furan (Berry, J. M.; Watson, C. Y.; Whish, W. J. D.; Threadgill, M. D. *J. Chem. Soc. Perkin Trans. 1* **1997**, 8, 1147) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.2 (M+H)⁺.

Example 211Preparation of 3-(2-isopropyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-methyl-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 345.2 (M+H)⁺.

Example 212

Preparation of 3-(2-isopropyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in
5 step 1(a) and 2-chloromethyl-3-chloro-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 365.2 (M+H)⁺.

Example 213

Preparation of 3-(2-isopropyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in
15 step 1(a) and 2-chloromethyl-5-methyl-thiophene (Moradpour, A. *J. Chem. Soc. Perkin Trans. 1*, **1993**, 1, 7) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 345.2 (M+H)⁺.

Example 214

Preparation of 3-(2-isopropyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in
25 step 1(a) and 2-chloromethyl-5-chloro-thiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 365.2 (M+H)⁺.

Example 215

Preparation of 5-(5-(2-isopropylphenylamino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in
step 1(a) and 5-chloromethyl-furan-2-carboxylic acid ethyl ester was
35 substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 387.2 (M+H)⁺.

Example 216Preparation of 5-(5-(2-isopropyl amino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carbaldehyde (Sanda, K.; Rigal, L.; Delmas, M.; Gaset, A. *Synthesis* **1992**, 6, 541) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 343.0 (M+H)⁺.

Example 217Preparation of 3-(2-isopropyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-thiophene (Lamy, J.; Lavit, D.; Buu-Hoi, N. P. *J. Chem. Soc.* **1958**, 4202) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 331.2 (M+H)⁺.

Example 218Preparation of 3-(2-isopropyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-furan (Arena, G.; Cali, R.; Maccarone, E.; Passerini, A. *J. Chem. Soc. Perkin Trans. 2* **1993**, 10, 1941) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.2 (M+H)⁺.

Example 219Preparation of 3-(3-methyl-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a), the title compound was prepared as a white solid. MS (ESI) 297.2 (M+H)⁺.

Example 220Preparation of 3-(3-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-

chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 221

5 Preparation of 3-(3-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

10

Example 222

Preparation of 3-(3-methyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.2 (M+H)⁺.

15

Example 223

Preparation of 3-(3-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 275.2 (M+H)⁺.

20

Example 224

Preparation of 3-(3-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.2 (M+H)⁺.

25

30

Example 225

Preparation of 3-(3-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step

35

1(d), the title compound was prepared as a white solid. MS (ESI) 302.2 (M+H)⁺.

Example 226

5 Preparation of 3-(3-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 311.4 (M+H)⁺.

10

Example 227

Preparation of 3-(3-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 333.2 (M+H)⁺.

15

Example 228

Preparation of 3-(3-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 327.2 (M+H)⁺.

20

Example 229

Preparation of 3-(3-methyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chlorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 331.2 (M+H)⁺.

25

30

Example 230

Preparation of 3-(3-methyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 327.2 (M+H)⁺.

35

Example 231Preparation of 3-(3-methyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole

5 Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-methylenedioxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 341.2 (M+H)⁺.

10

Example 232Preparation of 3-(3-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

15 Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 298.2 (M+H)⁺.

Example 233Preparation of 3-(3-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole

20 Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-furan (Berry, J. M.; Watson, C. Y.; Whish, W. J. D.; Threadgill, M. D. *J. Chem. Soc. Perkin Trans. 1* 1997, 8, 1147) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS
25 (ESI) 287.2 (M+H)⁺.

Example 234Preparation of 3-(3-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole

30 Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-methyl-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* 1988, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS
35 (ESI) 317.2 (M+H)⁺.

Example 235Preparation of 3-(3-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-chloro-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 337.2 (M+H)⁺.

Example 236Preparation of 3-(3-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-methyl-thiophene (Moradpour, A. *J. Chem. Soc. Perkin Trans. I*, **1993**, 1, 7) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 317.2 (M+H)⁺.

Example 237Preparation of 3-(3-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-chloro-thiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 337.2 (M+H)⁺.

Example 238Preparation of 5-(5-(3-methylphenylamino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester

Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carboxylic acid ethyl ester was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 359.2 (M+H)⁺.

Example 239Preparation of 3-(3-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-bromo-thiophene (Clapp, R. C.; Clark, J. H; Vaughan, J. R.; English, J. P.; Anderson, G. W. *J. Am. Chem. Soc.* **1947**, *60*, 1549) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 381.0 (M+H)⁺.

Example 240Preparation of 5-(5-(3-methylphenylamino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde

Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carbaldehyde (Sanda, K.; Rigal, L.; Delmas, M.; Gaset, A. *Synthesis* **1992**, *6*, 541) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.2 (M+H)⁺.

Example 241Preparation of 3-(3-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-thiophene (Lamy, J.; Lavit, D.; Buu-Hoi, N. P. *J. Chem. Soc.* **1958**, 4202) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 242Preparation of 3-(3-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-furan (Arena, G.; Cali, R.; Maccarone, E.; Passerini, A. *J. Chem. Soc. Perkin Trans. 2* **1993**, *10*, 1941) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 287.2 (M+H)⁺.

Example 243Preparation of 3-(4-*n*-butyl-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 339.2 (M+H)⁺.

Example 244Preparation of 3-(4-*n*-butyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 345.2 (M+H)⁺.

Example 245Preparation of 3-(4-*n*-butyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 357.2 (M+H)⁺.

Example 246Preparation of 3-(4-*n*-butyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 375.2 (M+H)⁺.

Example 247Preparation of 3-(4-*n*-butyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 317.2 (M+H)⁺.

Example 248Preparation of 3-(4-*n*-butyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 357.2 (M+H)⁺.

5

Example 249

Preparation of 3-(4-*n*-butyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 353.2 (M+H)⁺.

10

Example 250

Preparation of 3-(4-*n*-butyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chlorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 373.2 (M+H)⁺.

15

Example 251

Preparation of 3-(4-*n*-butyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 369.2 (M+H)⁺.

20

Example 252

Preparation of 3-(4-*n*-butyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-methylenedioxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 383.2 (M+H)⁺.

30

35

Example 253Preparation of 3-(4-*n*-butyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 344.2 (M+H)⁺.

Example 254Preparation of 3-(4-*n*-butyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 340.2 (M+H)⁺.

Example 255Preparation of 3-(4-*n*-butyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 369.2 (M+H)⁺.

Example 256Preparation of 3-(2,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 343.0 (M+H)⁺.

Example 257Preparation of 3-(2,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1 (d), the title compound was prepared as a white solid. MS (ESI) 349.0 (M+H)⁺.

Example 258Preparation of 3-(2,4-dimethoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole

5 Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 361.0 (M+H)⁺.

10

Example 259Preparation of 3-(2,4-dimethoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

15 Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 349.0 (M+H)⁺.

20

Example 260Preparation of 3-(2,4-dimethoxy-anilino)- (3,4-difluoro-benzylthio)-1,2,4-triazole

25 Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 379.0 (M+H)⁺.

Example 261

30 Preparation of 3-(2,4-dimethoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

35 Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 321.2 (M+H)⁺.

Example 262Preparation of 3-(2,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 361.0 (M+H)⁺.

Example 263Preparation of 3-(2,4-dimethoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 357.2 (M+H)⁺.

Example 264Preparation of 3-(2,4-dimethoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chlorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 377.0 (M+H)⁺.

Example 265Preparation of 3-(2,4-dimethoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 373.2 (M+H)⁺.

Example 266Preparation of 3-(2,4-dimethoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-methylenedioxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 387.2 (M+H)⁺.

Example 267Preparation of 3-(2,4-dimethoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 348.0 (M+H)⁺.

Example 268Preparation of 3-(2,4-dimethoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 344.2 (M+H)⁺.

Example 269Preparation of 3-(2,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 373.2 (M+H)⁺.

Example 270Preparation of 3-(2-methyl-4-methoxy-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 327.2 (M+H)⁺.

Example 271Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1 (d), the title compound was prepared as a white solid. MS (ESI) 333.2 (M+H)⁺.

Example 272Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 345.0 (M+H)⁺.

Example 273Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 333.2 (M+H)⁺.

Example 274Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 363.0 (M+H)⁺.

Example 275

Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 305.2 (M+H)⁺.

Example 276

Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 345.0 (M+H)⁺.

Example 277

Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 341.2 (M+H)⁺.

Example 278

Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chlorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 361.0 (M+H)⁺.

Example 279

Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 357.2 (M+H)⁺.

Example 280

Preparation 3-(2-methyl-4-methoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-methylenedioxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 371.0 (M+H)⁺.

Example 281

Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 332.2 (M+H)⁺.

Example 282

Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 328.2 (M+H)⁺.

Example 283

Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 357.2 (M+H)⁺.

Example 284

Preparation of 3-(2,6-dimethyl-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,6-dimethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 311.4 (M+H)⁺.

Example 285

Preparation of 3-(2,6-dimethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,6-dimethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 329.2 (M+H)⁺.

Example 286

Preparation of 3-(2,6-dimethyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,6-dimethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in

step 1(d), the title compound was prepared as a white solid . MS (ESI) 317.2 (M+H)⁺.

Example 287

5 Preparation of 3-(2,6-dimethyl-anilino)- (3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,6-dimethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide
10 in step 1(d), the title compound was prepared as a white solid. MS (ESI) 347.0 (M+H)⁺.

Example 288

15 Preparation of 3-(2,6-dimethyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,6-dimethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI)
20 289.0 (M+H)⁺.

Example 289

Preparation of 3-(2,6-dimethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,6-dimethylphenyl isothiocyanate was substituted for phenylisothiocyanate in
25 step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 329.2 (M+H)⁺.

Example 290

30 Preparation of 3-(2,6-dimethyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,6-dimethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in
35 step 1(d), the title compound was prepared as a white solid. MS (ESI) 325.2 (M+H)⁺.

Example 291Preparation of 3-(2,6-dimethyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,6-dimethylphenyl isothiocyanate was substituted for phenylisothiocyanate in
5 step 1(a) and 2-chlorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 345.0 (M+H)⁺.

Example 29210 Preparation of 3-(4-fluoro-anilino)-5-(furan-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-furan (Berry, J. M.; Watson, C. Y.; Whish, W. J. D.; Threadgill, M. D. *J. Chem. Soc. Perkin Trans. 1* 1997, 8, 1147) was substituted for benzyl
15 bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 291.2 (M+H)⁺.

Example 29320 Preparation of 3-(4-fluoro-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-methyl-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* 1988, 29(1), 117) was substituted for benzyl
25 bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 321.0 (M+H)⁺.

Example 29430 Preparation of 3-(4-fluoro-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-chloro-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* 1988, 29(1), 117) was substituted for benzyl
35 bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 341.2 (M+H)⁺.

Example 295Preparation of 3-(4-fluoro-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl
5 isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-methyl-thiophene (Moradpour, A. *J. Chem. Soc. Perkin Trans. 1*, 1993, 1, 7) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 321.0 (M+H)⁺.

Example 296Preparation of 3-(4-fluoro-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl
15 isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-chloro-thiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 341.2 (M+H)⁺.

Example 297Preparation of 5-(5-(4-fluorophenylamino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl
isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carboxylic acid ethyl ester was substituted for benzyl
25 bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 363.2 (M+H)⁺.

Example 298Preparation of 3-(4-fluoro-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl
isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-bromo-thiophene (Clapp, R. C.; Clark, J. H.; Vaughan, J. R.; English, J. P.; Anderson, G. W. *J. Am. Chem. Soc.* 1947, 60, 1549) was
35 substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 385.0 (M)⁺.

Example 299Preparation of 5-(5-(4-fluorophenylamino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carbaldehyde (Sanda, K.; Rigal, L.; Delmas, M.; Gaset, A. *Synthesis* **1992**, 6, 541) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 319.0 (M+H)⁺.

Example 300Preparation of 3-(4-fluoro-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-thiophene (Lamy, J.; Lavit, D.; Buu-Hoi, N. P. *J. Chem. Soc.* **1958**, 4202) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 307.2 (M+H)⁺.

Example 301Preparation of 3-(4-fluoro-anilino)-5-(furan-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-furan (Arena, G.; Cali, R.; Maccarone, E.; Passerini, A. *J. Chem. Soc. Perkin Trans. 2* **1993**, 10, 1941) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 291.2 (M+H)⁺.

Example 302Preparation of 3-methyl-3-anilino-5-benzylthio-1,2,4-triazolea) 3-anilino-5-benzylthio-1 or/2-methyl ethyl ether-1,2,4-triazole

To a stirring solution of 3-anilino-5-benzylthio-1,2,4-triazole (0.68 g, 2.41 mmol) in 8 mL DMF was added NaH (0.125 g, 3.13 mmol). To this mixture was added chloromethyl ethyl ether (0.251 g, 2.65 mmol), and the solution was stirred overnight. The reaction mixture was poured into 50 ml H₂O and extracted three times with EtOAc. The EtOAc extracts were dried over Na₂SO₄, filtered, and concentrated down. The crude mixture was subjected to column chromatography (silica gel, EtOAc/hexane) to provide the title compounds as a mixture of regioisomers as a light yellow oil (0.58 g,

71%). ¹H-NMR (400MHz, d6-DMSO) compound 1: δ9.33 (broad singlet, 1H), 7.51 (d, 2H, J=8.3 Hz), 7.42-7.22 (m, 8H), 5.23 (s, 2H), 4.47 (s, 2H), 3.43 (q, 2H, J=7.2 Hz), 1.04 (t, 3H, J=7.0 Hz). Compound 2: δ9.20 (broad singlet, 1H), 7.63 (d, 2H, J=7.6 Hz), 7.42-6.93 (m, 8H), 5.44 (s, 2H), 4.30 (s, 2H), 3.51 (q, 2H, J=7.1 Hz), 1.07 (t, 3H, J=7.0). MS (ESI) 341 (M+H)⁺.

b) 3-methyl-3-anilino-5-benzylthio-1,2,4-triazole

To a stirring solution of 3-anilino-5-benzylthio-1 or/2-methyl ethyl ether-1,2,4-triazole (50 mg, 0.15 mmol) in 1 ml THF was added NaH (11.8 mg, 0.30 mmol), and to this solution was added CH₃I (0.036 ml, 0.57 mmol). The reaction mixture was stirred overnight. THF was removed and 0.5 ml TFA was added to the residue and stirred overnight. TFA was removed under vacuum and the mixture was purified by preparative HPLC to afford the title compound as a clear oil (28 mg, 53%). ¹H-NMR (400MHz, d6-DMSO) δ3.3-7.25 (m, 10H), 4.27 (s, 2H), 3.40 (s, 3H). MS (ESI) 297 (M+H)⁺.

Example 303

Preparation of 3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except iodoethane was used in step 302(b) instead of iodomethane, the title compound was isolated as a white solid. ¹H-NMR (400MHz, d6-DMSO) δ7.42-7.26 (m, 10H), 4.26 (s, 2H), 3.86 (m, 2H), 1.20 (m, 3H). MS (ESI) 311 (M+H)⁺.

Example 304

Preparation of 3-n-propyl-3-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except 1-iodopropane was used in step 302(b) instead of iodomethane, the title compound was isolated as a white solid (35%). ¹H-NMR (400MHz, d6-DMSO) δ7.42-7.26 (m, 10H), 4.25 (s, 2H), 3.76 (t, 2H, J=6.5 Hz), 3.31 (t, 2H, J=1.4 Hz), 1.63 (m, 2H), 0.93 (t, 3H, J=7.4 Hz). MS (ESI) 325 (M+H)⁺.

Example 305

Preparation of 3-n-butyl-3-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except 1-iodobutane was used in step 302(b) instead of iodomethane, the title compound was isolated as a white solid (31%). ¹H-NMR (400MHz, d6-DMSO) δ7.42-7.22 (m, 10H), 4.26 (s, 2H), 3.80 (t, 2H, J=7.5 Hz), 3.31 (t, 2H, J=1.4 Hz), 1.59 (m, 2H), 1.36 (m, 2H), 0.92 (t, 3H, J=7.3 Hz). MS (ESI) 338 (M+H)⁺.

Example 306Preparation of 3-isopropyl-3-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except 1-iodo-2-methyl propane was used in step 302(b) instead of iodomethane, the title compound was isolated as a white solid. ¹H-NMR (400MHz, d6-DMSO) δ7.42-7.22 (m, 10H), 4.25 (s, 2H), 3.66 (d, 2H, J=7.6 Hz), 1.92 (m, 1H), 0.93 (d, 6H, J=6.7 Hz). MS (ESI) 338 (M+H)⁺.

Example 307Preparation of 3-allyl-3-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except allyl bromide was used in step 302(b) instead of iodomethane, the title compound was isolated as a white solid (41%). ¹H-NMR (400MHz, d6-DMSO) δ7.37-7.28 (m, 10H), 5.96 (m, 1H), 5.18 (m, 2H), 4.45 (s, 2H), 4.26 (s, 2H). MS (ESI) 323 (M+H)⁺.

Example 308Preparation of 3-benzyl-3-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except benzyl bromide was used in step 302(b) instead of iodomethane, the title compound was isolated as a white solid (48%). ¹H-NMR (400MHz, d6-DMSO) δ7.28-7.23 (m, 15H), 5.09 (s, 2H), 4.26 (s, 2H). MS (ESI) 373 (M+H)⁺.

Example 309Preparation of 3-methylacetate-3-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except methyl bromoacetate was used in step 302(b) instead of iodomethane, the title compound was isolated as a white solid. ¹H-NMR (400MHz, d6-DMSO) δ7.37-7.22 (m, 10H), 4.59 (s, 2H), 4.26 (s, 2H), 3.74 (s, 3H). MS (ESI) 355 (M+H)⁺.

Example 310Preparation of 3-methylacetate-3-(p-methyl)-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except 3-(p-methyl)-anilino-5-benzylthio-1,2,4-triazole was used in step 302(a) instead of 3-

anilino-5-benzylthio-1,2,4-triazole and methyl bromoacetate was used in step 302(b) instead of iodomethane, the title compound was isolated as a clear oil. ¹H-NMR (400MHz, d6-DMSO) δ7.38-7.09 (m, 9H), 4.56 (s, 2H), 4.27 (s, 2H), 3.75 (s, 3H), 2.37 (s, 3H). MS (ESI) 369 (M+H)⁺.

5

Example 311

Preparation of 3-methylacetate-3-(p-methoxy)-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except 3-(p-methoxy)-anilino-5-benzylthio-1,2,4-triazole was used in step 302(a) instead of 3-anilino-5-benzylthio-1,2,4-triazole and methyl bromoacetate was used in step 302(b) instead of iodomethane, the title compound was isolated as a brown oil (44%). ¹H-NMR (400MHz, d6-DMSO) δ7.92-7.22 (m, 7H), 6.99 (d, 2H, J=8.9 Hz), 4.51 (s, 2H), 4.26 (s, 2H), 3.83 (s, 3H), 3.76 (s, 3H). MS (ESI) 385 (M+H)⁺.

15

Example 312

Preparation of 3-methylacetate-3-(2,6-dimethyl)-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except 3-(2,6-dimethyl)-anilino-5-benzylthio-1,2,4-triazole was used in step 302(a) instead of 3-anilino-5-benzylthio-1,2,4-triazole and methyl bromoacetate was used in step 302(b) instead of iodomethane, the title compound was isolated as a white solid (43%). ¹H-NMR (400MHz, d6-DMSO) δ7.32-7.19 (m, 8H), 4.37 (s, 2H), 4.25 (s, 2H), 3.77 (s, 3H), 2.27 (s, 6H). MS (ESI) 383 (M+H)⁺.

25

Biological Data:

Direct Spectrophotometric Assays of hMetAP2:




The hMetAP2 activity can be measured by direct spectrophotometric assay methods using alternative substrates, L-methionine-p-nitroanilide (Met-pNA) and L-methionine-7-amido-4-methylcoumarin (Met-AMC). The formation of p-nitroaniline (pNA) or 7-amido-4-methylcoumarin (AMC) was continuously monitored by increasing absorbance or fluorescence at 405 nm and 460 nm, respectively, on a corresponding plate reader. All assays were carried out at 30°C. The fluorescence or spectrophotometric plate reader was calibrated using authentic pNA and AMC from Sigma, respectively. For a typical 96-well plate assay, the increase in the absorbance (at 405 nm for pNA) or the fluorescence

35

emission ($\lambda_{\text{ex}} = 360 \text{ nm}$, $\lambda_{\text{em}} = 460 \text{ nm}$, for AMC) of a 50 μL assay solution in each well was used to calculate the initial velocity of hMetAP2. Each 50 μL assay solution, contained 50 mM Hepes- Na^+ (pH 7.5), 100 mM NaCl, 10-100nM purified hMetAP2 enzyme, and varying amounts of Met-AMC (in 3% DMSO aqueous solution) or Met-pNA. Assays were initiated with the addition of substrate and the initial rates were corrected for the background rate determined in the absence of hMetAP2.

Coupled Spectrophotometric Assays of hMetAP2:

The methionine aminopeptidase activity of hMetAP2 can also be measured spectrophotometrically by monitoring the free L-amino acid formation. The release of N-terminal methionine from a tripeptide (Met-Ala-Ser, Sigma) or a tetrapeptide (Met-Gly-Met-Met, Sigma) substrate was assayed using the L-amino acid oxidase (AAO) / horse radish peroxidase (HRP) couple (eq. 1-3a,b). The formation of hydrogen peroxide (H_2O_2) was continuously monitored at 450nm (absorbance increase of o-Dianisidine (Sigma) upon oxidation, $\Delta\epsilon = 15,300 \text{ M}^{-1}\text{cm}^{-1}$)² and 30 °C in a 96- or 384-well plate reader by a method adapted from Tsunasawa, S. et al.(1997) (eq. 3a). Alternatively, formation of H_2O_2 was followed by monitoring the fluorescence emission increase at 587nm ($\Delta\epsilon = 54,000 \text{ M}^{-1}\text{cm}^{-1}$, $\lambda_{\text{ex}} = 563 \text{ nm}$, slit width for both excitation and emission was 1.25 mm) and 30 °C using Amplex Red (Molecular Probes, Inc) (Zhou, M. et al. (1997) *Anal. Biochem.* 253, 162) (eq. 3b). In a total volume of 50 μL , a typical assay contained 50 mM Hepes- Na^+ , pH 7.5, 100 mM NaCl, 10 μM CoCl_2 , 1 mM o-Dianisidine or 50 μM Amplex Red, 0.5 units of HRP (Sigma), 0.035 unit of AAO (Sigma), 1 nM hMetAP2, and varying amounts of peptide substrates. Assays were initiated by the addition of hMetAP2 enzyme, and the rates were corrected for the background rate determined in the absence of hMetAP2.



Kinetic Data Analysis:

Data were fitted to the appropriate rate equations using Grafit computer software. Initial velocity data conforming to Michaelis-Menton kinetics were fitted to eq. 4. Inhibition patterns conforming to apparent competitive and non-competitive inhibition were fitted to eq. 5 and eq. 6, respectively.

$$v = VA/(K_a + A) \quad (4)$$

$$v = VA/[K_a(1 + I/K_{is}) + A] \quad (5)$$

$$v = VA/[K_a(1 + I/K_{is}) + A(1 + I/K_{ii})] \quad (6)$$

In eqs. 4 - 6, v is the initial velocity, V is the maximum velocity, K_a is the apparent Michaelis constant, I is the inhibitor concentration, and A is the concentration of variable substrates. The nomenclature used in the rate equations for inhibition constants is that of Cleland (1963), in which K_{is} and K_{ii} represent the apparent slope and intercept inhibition constants, respectively.

Cell growth inhibition assays:

The ability of MetAP2 inhibitors to inhibit cell growth was assessed by the standard XTT microtitre assay. XTT, a dye sensitive to the pH change of mitochondria in eukaryotic cells, is used to quantify the viability of cells in the presence of chemical compounds. Cells seeded at a given number undergo approximately two divisions on average in the 72 hours of incubation. In the absence of any compound, this population of cells is in exponential growth at the end of the incubation period; the mitochondrial activity of these cells is reflected in the spectrophotometric readout (A_{450}). Viability of a similar cell

population in the presence of a given concentration of compound is assessed by comparing the A450 reading from the test well with that of the control well. Flat-bottomed 96-well plates are seeded with appropriate numbers of cells ($4-6 \times 10^3$ cells/well in a volume of 200 μ l) from trypsinized exponentially growing cultures. In the case of HUVECs, the wells are coated with matrigel prior to establishing the cultures. To "blank" wells is added growth medium only. Cells are incubated overnight to permit attachment. Next day, medium from wells that contain cells is replaced with 180 μ l of fresh medium. Appropriate dilutions of test compounds are added to the wells, final DMSO concentration in all wells being 0.2 %. Cells plus compound are incubated for an additional 72 hr at 37°C under the normal growth conditions of the cell line used. Cells are then assayed for viability using standard XTT/PMS (prepared immediately before use: 8 mg XTT (Sigma X-4251) per plate is dissolved in 100 μ l DMSO. 3.9 ml H₂O is added to dissolve XTT and 20 μ l of PMS stock solution (30 mg/ml) is added from frozen aliquoted stock solution (10 mg of PMS (phenazine methosulfate, Sigma P-9625) in 3.3 ml PBS without cations. These stocks are frozen at -20°C until use). 50 μ l of XTT/PMS solution is added to each well and plates incubated for 90 minutes (time required may vary according to cell line, etc.) at 37°C until A₄₅₀ is >1.0. Absorbance at 450 nm is determined using a 96-well UV plate reader. Percent viability of cells in each well is calculated from these data (having been corrected for background absorbance). IC₅₀ is that concentration of compound that reduces cell viability to 50% control (untreated) viability.

The compounds of this invention show MetAP2 inhibitor activity having IC₅₀ values in the range of 0.0001 to 100 μ M. The full structure/activity relationship has not yet been established for the compounds of this invention. However, given the disclosure herein, one of ordinary skill in the art can utilize the present assays in order to determine which compounds of this invention are inhibitors of MetAP2 and which bind thereto with an IC₅₀ value in the range of 0.0001 to 100 μ M.

All publications, including, but not limited to, patents and patent applications cited in this specification, are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

The above description fully discloses the invention including preferred embodiments thereof. Modifications and improvements of the embodiments

- specifically disclosed herein are within the scope of the following claims. Without further elaboration it is believed that one skilled in the art can, given the preceding description, utilize the present invention to its fullest extent. Therefore any examples are to be construed as merely illustrative and not a limitation on the scope of the
- 5 present invention in any way. The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215